Due: September 15th

## Problem 1 (10 pts)

Consider a tensor  $T = \sum_{i=1}^r u_i \otimes v_i \otimes w_i \in \mathbb{R}^{d_1 \times d_2 \times d_3}$ . Recall that  $T_a := \sum_{i=1}^r (w_i)_a u_i \otimes v_i$ . Prove that for all  $a \in [d_3]$ , we have

- $\operatorname{colspan}(T_a) \subseteq \operatorname{span}(\{u_i\})$
- $\operatorname{rowspan}(T_a) \subseteq \operatorname{span}(\{v_i\})$

## Problem 2 (5 pts)

Show that for a matrix M, its rank and its border rank are always the same. In particular, suppose you are given a matrix M and a parameter r so that for every  $\epsilon > 0$  there is a rank r matrix  $M_r$  so that M and  $M_r$  are entrywise  $\epsilon$ -close. Show that rankM is the Eckhart-Young Theorem.

## Problem 3 (45 pts)

Let  $T = \sum_{i=1}^{d} \lambda_i u_i^{\otimes 3}$  for  $u_1, \dots, u_d \in \mathbb{R}^d$  a collection of orthonormal vectors and  $\lambda_1, \dots, \lambda_d > 0$ . Let  $p : \mathbb{R}^d \to \mathbb{R}$  be the associated cubic polynomial

$$p(x) := \sum_{i=1}^{d} \lambda_i \langle u_i, x \rangle^3.$$

The result we will prove is the following.

**Theorem 1.** Let  $x \in \mathbb{S}^{d-1}$  be any point for which  $p(x) \geq 0$ . Then x is a strict local maximum<sup>1</sup> for p over the domain  $\mathbb{S}^{d-1}$  if and only if  $x = u_i$ .

In other words, there are no "spurious local maxima" for the objective function p, which means that we can find a component  $u_i$  of the tensor T simply by running an appropriate implementation of gradient descent.

1. (2 pts) It suffices to establish Theorem 1 when  $u_1, \ldots, u_d$  are the standard basis vectors, i.e. when  $u_i = (0, \ldots, 0, 1, 0, \ldots, 0)$  where the 1 entry is in the *i*-th coordinate. Give an informal argument for why this is the case.

For the remaining sub-problems, we assume that  $\{u_i\}$  are the standard basis vectors.

2. (10 pts) Prove that every  $u_i$  is a strict local maximum.

We say a point  $x \in \mathbb{S}^{d-1}$  is a strict local maximum for a function  $f: \mathbb{S}^{d-1} \to \mathbb{R}$  if there exists some  $\epsilon > 0$  such that for all  $x' \in \mathbb{S}^{d-1}$  which are distinct from x and which satisfy  $||x' - x|| \le \epsilon$ , we have f(x') < f(x).

- 3. (2 pts) Prove that any local maximum x of p over  $\mathbb{S}^{d-1}$  is a stationary point, that is, the projection of the gradient of p at x to the tangent space at x is zero.
- 4. (3 pts) Prove that any local maximum x satisfies

$$\lambda_i x_i^2 = p(x) \cdot x_i \quad \text{for all } i \in [d].$$
 (1)

(Here  $x_i$  denotes the *i*-th coordinate of x.)

5. (3 pts) Prove that if  $x \in \mathbb{S}^{d-1}$  satisfies p(x) = 0, then x is not a strict local maximum.

Now we arrive at the trickiest part of the proof. Let  $x \in \mathbb{S}^{d-1}$  be any point that satisfies p(x) > 0 and has more than one nonzero coordinate. We need to show that such an x cannot be a local maximum. Let  $S^* \subseteq [d]$  denote the subset of coordinates of x which are nonzero.

- 6. **(5 pts)** Let  $S \subseteq S^*$  be any proper subset of  $S^*$ . Construct a unit vector w such that (1)  $\langle w, x \rangle = 0$ , (2) the entries of w indexed by S are given by scaling all the entries of x indexed by S by the same factor, and (3) the entries of w indexed by S are given by scaling all the entries of S indexed by S by the same factor.
- 7. (20 pts) Prove that x is not a strict local maximum.

  Hint: Perturb x in the direction of w and rescale it to give a unit vector  $x' = \delta w + \sqrt{1 \delta^2} x$ . Argue that p achieves a higher value at x' than at x. You may find Eq. (1) helpful.